

# PLASMA PHYSICS GROUP

*N72-29208*

## Current-Driven Electrostatic and Electromagnetic Ion Cyclotron Instabilities

D. W. Forslund

C. F. Kennel


J. M. Kindel

February 1971

R-87

CASE FILE  
COPY

DEPARTMENT OF PHYSICS  
UNIVERSITY OF CALIFORNIA  
LOS ANGELES 90024



Current-Driven Electrostatic and Electromagnetic  
Ion Cyclotron Instabilities

D. W. Forslund

C. F. Kennel

J. M. Kindel

February 1971

R-87

Plasma Physics Group  
Department of Physics  
University of California  
Los Angeles, California 90024

This work was partially supported by the Office of Naval Research, Grant #N00014-69-A-0200-4023; the National Science Foundation, Grant #GP-22817; the Atomic Energy Commission, Contract AT(04-3)-34, Project #157; and the National Aeronautics and Space Administration, Contract #NGL-05-007-190.

# Current-Driven Electrostatic and Electromagnetic Ion Cyclotron Instabilities

D. W. Forslund

Los Alamos Scientific Laboratory, Los Alamos, New Mexico 87544

C. F. Kennel\*

Department of Physics, UCLA, Los Angeles, California 90024

and

J. M. Kindel

Plasma Physics Laboratory, Princeton University,  
Princeton, New Jersey 08540

## ABSTRACT

Growth rates and parameter dependences are calculated for the current-driven instabilities of electrostatic (with finite- $\beta$  corrections) and electromagnetic ion cyclotron waves. For  $0.25 < T_e/T_i < 2.5$ , ion cyclotron waves have large growth rates, while ion acoustic waves are still stable. In fusion devices, where electrostatic waves may be stable, electromagnetic ion cyclotron waves are unstable for  $\beta_i > 10^{-3}$ . The suggestion is made that ion cyclotron waves may provide an anomalous resistance, with

$$\nu_{\text{eff}} \approx A^{1/2} \Omega_{ci} (T_{e,\parallel}/T_{i,\perp}).$$

## INTRODUCTION

Ion cyclotron instabilities, driven by magnetic-field-aligned currents and/or electron heat conduction, are of interest to the physics of the solar wind,<sup>1</sup> the ionosphere,<sup>2</sup> Q-machine plasmas,<sup>3-5</sup> and to certain controlled thermonuclear fusion devices such as tokamaks, which have confining electron currents. We report recent salient results concerning four basic extensions of the Drummond-Rosenbluth<sup>6</sup> investigation of the electrostatic ion cyclotron instability: arbitrary  $T_e/T_i$ , where  $T_e/T_i$  is the electron-to-ion temperature ratio; growth rates well above marginal stability; effects of contaminations by low charge-to-mass ratio ions; finite  $\beta_i$  corrections, where  $\beta_i$  is the ratio of ion particle to magnetic field pressure. Moreover, we report some preliminary results concerning the electromagnetic ion cyclotron current instability,<sup>7,8</sup> and discuss a few implications of saturated ion cyclotron turbulence. The details of this work will be published elsewhere.<sup>9</sup>

## ELECTROSTATIC ION CYCLOTRON INSTABILITIES

For purposes of analysis, we have chosen isotropic Maxwellian distributions (albeit with different  $T_e$  and  $T_i$ ) drifting along a uniform-background magnetic field  $B_0 \hat{e}_z$ . Thus, we have neglected the effect of the self-consistent magnetic field created by the current. To facilitate calculation in analytically difficult parameter regimes, a computer program has been developed which solves the dispersion relation for small-amplitude electrostatic waves.<sup>6,7</sup> The general electrostatic dispersion relation is cast into the Gordeyev integral form.<sup>10</sup> Complex roots of the dispersion relation are

obtained numerically by a standard algorithm, such as Muller's method.<sup>11</sup> A given wave mode is found either by making a suitably accurate guess or by following known wave modes into the parameter range of interest.

### SINGLE ION SPECIES

A marginal stability analysis<sup>2, 6, 8</sup> indicates that when  $T_e/T_i = 1$ , the wave that is marginally stable to the smallest current has  $K_\perp^2 R_i^2 \approx 1$ ,  $\omega \approx 1.2 \Omega_{ci}$ , and  $K_\perp/K_\parallel = 12$ , where  $R_i$  is the thermal ion Larmor radius,  $\omega$  is the wave frequency, and  $K_\perp/K_\parallel$  is the ratio of wave numbers perpendicular and parallel to  $B_0$ . Here, we focus on unstable ion cyclotron waves. The general electrostatic dispersion relation<sup>6, 7</sup> has dependences upon the following parameters:  $K_\perp^2 R_i^2$ ,  $K_\perp/K_\parallel$ ,  $V_D/a_e$  [the electron drift relative to ions in units  $(2T_e/m_e)^{1/2}$ ],  $m/M$ ,  $T_e/T_i$ , and the ratio of ion plasma to cyclotron frequency  $\omega_{pi}/\Omega_{ci}$ . We can eliminate some dependences: We consider  $\omega_{pi}^2/\Omega_{ci}^2 > 10$ , since this implies weak  $\omega_{pi}^2/\Omega_{ci}^2$  dependence.<sup>9</sup> We eliminate the dependences upon  $K_\perp/K_\parallel$  and  $K_\perp^2 R_i^2$  by searching numerically for local maxima in the growth rate. Choosing  $M/m = 1836$ , corresponding to an  $H^+$  plasma, we plot in Fig. 1 contours of maximum growth rate for the  $n = 1$  wave as a function of the two remaining parameters,  $V_D/a_e$  and  $T_e/T_i$ . For comparison, we also plot the marginal stability contour for the  $K_\perp = 0$  ion acoustic wave, corresponding to the magnetic field-free calculations of Fried and Gould<sup>12</sup> and Stringer,<sup>13</sup> and the contour for  $\gamma_{\max}/\omega_{pi} = 1$ , which marks the transition to the nonresonant electron-ion beam instability originally studied by Buneman. The shaded region

denotes the portion of the  $T_e/T_i$ ,  $V_D/a_e$  parameter space for which the ion cyclotron wave is unstable but the ion acoustic wave is stable. The ion cyclotron wave is unstable to the smaller current over the range  $0.05 < T_e/T_i < 8.5$ . Furthermore, it has a significant growth rate  $\gamma_{\max}/\Omega_i = 0.1$  (while the ion acoustic wave is still stable) over the range  $0.25 < T_e/T_i < 2.5$ . Thus, the ion cyclotron instability dominates the ion acoustic instability over an interestingly large portion of parameter space. It can be shown that the region of ion cyclotron dominance increases as  $M_i$  increases.<sup>2</sup>

## CYCLOTRON INSTABILITIES IN MULTI-ION PLASMAS

We now turn briefly to the operation of the electrostatic ion cyclotron instability in ion mixtures. Besides the above-listed parameters, there is now an additional dependence on  $N_i/N_e$ , the relative ion composition. Only the marginal stability ( $\gamma_{\max}/\Omega_{ci} = 0$ ) contour is considered here because of the added dimensionality in parameter space. We may ask the following questions: First, in a mixture, the ion harmonic of which ion is unstable to the smallest current? Second, how do critical electron drifts respond to changes in ion composition? Third, at what point does the plasma switch ion instabilities as the ion composition is varied? fourth, and last, how do these effects vary with  $T_e/T_i$ ? Figure 2 shows the results of an analytic calculation<sup>2</sup> of the electron drift  $V_c$  for marginal stability in a mixture of  $H^+$  and  $O^+$  (a case of ionospheric interest) as a function of  $T_e/T_i$  and fractional  $O^+$  concentration,  $N_{O^+}/N_e$ . It is convenient to represent our results in  $N_{O^+}/N_e$ ,



$V_D/a_c$  space with  $T_e/T_i$  as a parameter. Solid and dashed lines denote the marginal stability curves for the instabilities near the first  $O^+$  and  $H^+$  cyclotron harmonics, respectively. The appropriate temperature ratio  $T_e/T_i$  is indicated at the intersections of the  $H^+$  and  $O^+$  curves. For simplicity, the  $H^+$  and  $O^+$  temperatures were taken equal. Figure 2 indicates that the heavy-ion harmonic is unstable to the smaller current for surprisingly small heavy-ion contaminations, of the order of 10%.

### ELECTROMAGNETIC ION CYCLOTRON WAVES

A computer program which solves the full electromagnetic wave dispersion relation for spatially homogeneous, drifting electron and ion Maxwellian distributions in a uniform magnetic field has also been developed. In addition to the single ion parameters listed above, the dispersion relation now depends upon  $\beta_i$ . For large-enough  $\beta_i$ , it has been shown by Stix<sup>7</sup> that electromagnetic ion cyclotron waves (which have electrostatic corrections at oblique propagation) have a parallel phase velocity sufficiently slow that they can be destabilized by small electron drifts. Details of our investigation of this instability will be considered elsewhere.<sup>9</sup>

Figure 3 shows some of our preliminary results. For  $M/m = 1836$  and  $T_e/T_i = 1$ , we plot contours of constant maximum growth rate in a  $\beta_i$ ,  $V_D/a_e$  parameter space.

For purposes of comparison, we have also plotted the marginal stability contour,  $\gamma_{\max}/\Omega_i = 0$ , for the "electrostatic" ion cyclotron wave. This mode is electrostatically polarized for  $\beta_i$  small, and develops small

electromagnetic components as  $\beta_i$  increases. These finite  $\beta_i$  corrections are stabilizing. However, ion cyclotron waves of one sort or another continue to be unstable as  $\beta_i$  increases, because of the lower threshold of this new electromagnetic instability for  $\beta_i > 10^{-3}$ . It is probably this mode which is most important in the solar wind, since  $\beta_i > 10^{-2}$ , even near the sun.

For  $\gamma_{\max}/\Omega_i = 10^{-2}$ , the real frequency is typically  $0.75\Omega_i$ , and  $K_{\perp}R_i \approx 1.0 - 1.5$ ,  $K_{\parallel}R_i < 0.1$  for maximum growth. As the drift velocity is decreased to the point where  $\gamma_{\max}/\Omega_i < 10^{-3}$ ,  $K_{\perp}R_i$  for maximum growth drops rapidly below 1.0 and  $K_{\parallel}R_i$  increases somewhat above 0.1. Drifts then tend toward Stix's estimate.

Before closing the discussion of the electromagnetic current instability, let us note that a drifting electron Maxwellian certainly is not the self-consistent equilibrium distribution in a finite  $\beta_i$  current-carrying plasma. However, the Maxwellian should reasonably well describe the electron growth. Furthermore, any stabilizing effect of the inhomogeneous magnetic field, created by the direct current, might be unimportant because of the small wave group velocity. Kan has indicated to us that if the inhomogeneous magnetic field determines a pressure gradient, a drift mode might also be easily destabilized for  $\beta_i > 10^{-2}$ .

#### NONLINEAR CONSIDERATIONS

One-dimensional quasilinear electron plateau formation can saturate the electrostatic ion cyclotron instability at  $e\phi/T_e \leq 10^{-2}$ ,  $\phi$  being the wave



potential integrated over the unstable spectrum.<sup>6,14</sup> Dum and Dupree,<sup>4</sup> however, have shown that if electron plateau formation is inhibited,  $e\phi/T_e \sim 1/2$ . This large saturation level agrees with Q machine results,<sup>3-5,15</sup> where end effects may prevent plateau formation. We have observed<sup>16</sup> that large  $e\phi/T_e$  permits ion cyclotron waves to interact with a significant portion of the electron distribution, at an effective electron-ion frequency  $\nu_{\text{eff}} \simeq A\Omega_{ci} (T_{e,\parallel}/T_{i,\perp})$ , where  $A$  is the atomic number, and  $\perp$  and  $\parallel$  refer to  $B_0$ . For large  $e\phi/T_e$ , anomalous resistance due directly to wave-electron scattering may occur, or turbulence may hold back the bulk of the electron distribution while high-energy electrons carry current demanded above cyclotron threshold. In the latter case, anomalous resistance ultimately may be derived from two-stream instabilities generated by the electron tail. Significant spatial plasma diffusion and perpendicular ion heating may also occur.

The conclusions described above could be important for tokamak controlled-fusion devices. Experiments in the Princeton Model ST may be operating in regimes where the electrostatic ion cyclotron wave is likely to be unstable, but where the acoustic wave may or may not be unstable.<sup>17-19</sup> It has been suggested that larger  $\beta_i$  (higher density) may eliminate electrostatic current instabilities by reducing the electron drift needed to produce the confining magnetic field. However, we must still contend with the electromagnetic ion cyclotron instability. Bers, Manheimer, and Coppi have already suggested that ion cyclotron waves may account for the anomalous perpendicular ion heating observed in the Soviet TM-3 Tokamak.<sup>20</sup>

## ACKNOWLEDGMENTS

\* Consultant to Los Alamos Scientific Laboratory.

This work was supported at Princeton by U. S. Atomic Energy Commission Contract AT(30-1)-1238; at UCLA by U. S. Atomic Energy Commission Contract AT(04-3)-34, Project No. 157, and NASA Grant NGR 05-007-190; at Los Alamos by U. S. Atomic Energy Commission Contract W-7405-ENG.36. C. F. Kennel was supported by an Alfred P. Sloan faculty fellowship.

## REFERENCES

- <sup>1</sup>D. W. Forslund, J. Geophys. Res. 75, 17 (1970).
- <sup>2</sup>J. M. Kindel and C. F. Kennel, J. Geophys. Res. (to be published).
- <sup>3</sup>N. Booth, D. R. Baker, R. L. Stenzel, and A. Y. Wong, Bull. Am. Phys. Soc. 15, 1460 (1970).
- <sup>4</sup>C. T. Dum and T. H. Dupree, Phys. Fluids 13, 2064 (1970).
- <sup>5</sup>T. K. Chu, H. W. Hendel, and A. Y. Wong, private communication.
- <sup>6</sup>W. E. Drummond and M. N. Rosenbluth, Phys. Fluids 5, 1507 (1962).
- <sup>7</sup>T. H. Stix, The Theory of Plasma Waves (McGraw-Hill, New York, 1962), pp. 202 and 223.
- <sup>8</sup>D. W. Forslund, C. F. Kennel, and J. M. Kindel, Bull. Am. Phys. Soc. 15, 1459 (1970).

- <sup>9</sup>D. W. Forslund, C. F. Kennel, and J. M. Kindel, private communication.
- <sup>10</sup>P. C. Clemmow and J. P. Dougherty, Electrodynamics of Particles and Plasmas (Addison-Wesley, Reading, Mass., 1969), p. 286.
- <sup>11</sup>R. D. Rodman, Commun. ACM 6, 442 (1963) and V. W. Whitley, Commun. ACM 11, 12 (1968).
- <sup>12</sup>B. D. Fried and R. W. Gould, Phys. Fluids 4, 139 (1961).
- <sup>13</sup>T. E. Stringer, J. Nucl. Energy, Pt. C 6, 267 (1965).
- <sup>14</sup>V. I. Petviashvili, Zh. Eksp. Teor. Fiz. 45, 1467 (1963)[Sov. Phys. - JETP 18, 1014 (1964)].
- <sup>15</sup>N. S. Buchel'nikova and R. A. Salimov, Zh. Eksp. Teor. Fiz. 56, 1108 (1969)[Sov. Phys. - JETP 29, 595 (1969)].
- <sup>16</sup>J. M. Kindel and C. F. Kennel, Bull. Am. Phys. Soc. 15, 1460 (1970).
- <sup>17</sup>D. J. Grove, D. L. Dimock, E. Hinnov, J. C. Hosea, L. C. Johnson, E. D. Meservy, and E. L. Tolnas, Princeton Plasma Physics Laboratory MATT-813 (1970).
- <sup>18</sup>F. W. Perkins and P. H. Rutherford, private communication.
- <sup>19</sup>A. Hirose, M. Murakami, and I. Alexeff, Oak Ridge National Laboratory Report ORNL-TM-2988 (1970).
- <sup>20</sup>A. Bers, W. Manheimer, and B. Coppi, Bull. Am. Phys. Soc. 15, 1419 (1970).

# FIGURE CAPTIONS

Fig. 1. Contours of maximum growth for the  $n = 1$   $H^+$  cyclotron wave (solid) and the  $H^+$  acoustic wave (dashed). The shaded region denotes where the cyclotron wave is unstable but the acoustic wave stable.

Fig. 2. Critical drift versus fractional  $O^+$  concentration for  $T_e/T_i = 1, 3, 5, 10$ . The solid (dashed) lines denotes the  $n = 1$ ,  $O^+$  ( $H^+$ ) critical drift. The transition from  $O^+$  to  $H^+$  waves shifts from  $N(O^+)/N_e = 0.15$ , at  $T_e/T_i = 1$ , to  $N(O^+)/N_e = 0.06$ , at  $T_e/T_i = 10$ .

Fig. 3. "Electrostatic" and electromagnetic ion cyclotron maximum growth rate contours for  $M/m = 1836$  and  $T_e/T_i = 1$ . The electrostatic mode has  $\text{Re}(\omega/\Omega_i) > 1$ , whereas  $\text{Re}(\omega/\Omega_i) < 1$  for the electromagnetic mode. The dashed extension occurs because of the difficulty of distinguishing the electrostatic from the electromagnetic modes. Above  $\beta_i \sim 10^{-3}$ , the electromagnetic mode is unstable to the smaller current.




Fig 1

712004




Fig 2

712005

UCLA PLASMA PHYSICS GROUP REPORTS

- R-1 "Propagation of Ion Acoustic Waves Along Cylindrical Plasma Columns", A.Y. Wong (July 1965)
- R-2 "Stability Limits for Longitudinal Waves in Ion Beam-Plasma Interaction", B.D. Fried and A.Y. Wong (August 1965)
- R-3 "The Kinetic Equation for an Unstable Plasma in Parallel Electric and Magnetic Fields", B.D. Fried and S.L. Ossakow (Nov. 1965)
- R-4 "Low-Frequency Spatial Response of a Collisional Electron Plasma", B.D. Fried, A.N. Kaufman and D.L. Sachs (Aug. 1965)
- R-5 "Effects of Collisions on Electrostatic Ion Cyclotron Waves", A.Y. Wong, O. Judd and F. Hui (Dec. 1965)
- R-6 "Interaction Between Ion Beams and Plasmas", R. Rowberg, A.Y. Wong and J.M. Sellen (April 1966)
- R-7 "Observation of Cyclotron Echoes from a Highly Ionized Plasma", D.E. Kaplan and R.M. Hill (May 1966)
- R-8 "Excitation and Damping of Drift Waves", A.Y. Wong and R. Rowberg (July 1966)
- R-9 "The Guiding Center Approximation in Lowest Order", Alfredo Banos, Jr. (Sept. 1966)
- R-10 "Plasma Streaming into a Magnetic Field", S.L. Ossakow (Nov. 1966)
- R-11 "Cooperative Effects in Plasma Echo Phenomena", A.Y. Wong (March 1967)
- R-12 "A Quantum Mechanical Study of the Electron Gas Via the Test Particle Method", M.E. Rensink (March 1967)
- R-13 "Linear and Nonlinear Theory of Grid Excitation of Low Frequency Waves in a Plasma", G.L. Johnston (April 1967)
- R-14 "The Expansion and Diffusion of an Isolated Plasma Column", J. Hyman (May 1967)
- R-15 "Two-pole Approximation for the Plasma Dispersion Function", B.D. Fried, C.L. Hedrick and J. McCune (August 1967)
- R-16 "Experimental Investigation of Electron Runaway Phenomena", J.S. DeGroot (August 1967)
- R-17 "Parametric Coupling Between Drift Waves", F. Hui, R. Rowberg and A.Y. Wong (Oct. 1967)
- R-18 "Cyclotron Echoes from Doppler Effects", A.Y. Wong (March 1968)
- R-19 "Ion Wave Echoes", D.R. Baker, N.R. Ahren and A.Y. Wong (Nov. 1967)
- R-20 "Cyclotron Echoes in Plasmas", O. Judd, Thesis (March 1968)
- R-21 "Test Particle Theory for Quantum Plasmas", M.E. Rensink (Oct. 1967)
- R-22 "Artificial Van Allen Belt", Charles F. Kennel (Nov. 1967)
- R-23 "Landau Damping of Ion Acoustic Waves in a Cesium Plasma with Variable Electron-Ion Temperature Ratio, K.B. Rajangam (Oct. 1967)
- R-24 "The Inhomogeneous Two-Stream Instability", G. Knorr (Sept. 1967)
- R-25 "Magnetic Turbulence in Shocks", C.F. Kennel and H.E. Petschek (Dec. 1967)
- R-26 "Small Amplitude in High Beta Plasmas", V. Formisano and C. Kennel (Feb. 1968)
- R-27 "Low Beta Plasma Penetration Across a Magnetic Field", B.D. Fried and S. Ossakow (March 1968)
- R-28 "Annual Status Report", Feb. 1, 1967-Jan. 31, 1968, Principal Investigators A. Banos, Jr., B.D. Fried, C.F. Kennel



- R-29 "The Theorist's Magnetosphere", C. Kennel (April 1968)
- R-30 "Electromagnetic Pitch Angle Instabilities in Space", C.F. Kennel and F.L. Scarf (April 1968)
- R-31 "Electromagnetic Echoes in Collisionless Plasmas," A.Y. Wong (April 1968)
- R-32 "Parametric Excitation of Drift Waves in a Resistive Plasma", G. Weyl and M. Goldman (June 1968)
- R-33 "Parametric Excitation from Thermal Fluctuations at Plasma Drift Wave Frequencies", A.Y. Wong, M.V. Goldman, F. Hai, R. Rowberg (May 1968)
- R-34 "Current Decay in a Streaming Plasma Due to Weak Turbulence ", S.L. Ossakow and E.D. Fried (June 1968)
- R-35 "Temperature Gradient Instabilities in Axisymmetric Systems", C.S.Liu (August 1968)
- R-36 "Electron Cyclotron Echo Phenomena in a Hot Collisionless Plasma", O. Judd (August 1968)
- R-37 "Transverse Plasma Wave Echoes", B.D. Fried and Craig Olson (Oct. 1968)
- R-38 "Low Frequency Interchange Instabilities of the Ring Current Belt", C.S. Liu (Jan. 1969)
- R-39 "Drift Waves in the Linear Regime", R.E. Rowberg and A.Y. Wong (Feb. 1969)
- R-40 "Parametric Mode-Mode Coupling Between Drift Waves in Plasmas", F. Hai and A.Y. Wong (Jan. 1969)
- R-41 "Nonlinear Oscillatory Phenomena with Drift Waves in Plasmas", F. Hai and A.Y. Wong (Sept. 1960)
- R-42 "Ion-Burst Excited by a Grid in a Plasma", H. Ikezi and R.J. Taylor (Feb. 1969)
- R-43 "Measurements of Diffusion in Velocity Space from Ion-Ion Collisions", A. Wong and D. Baker (March 1969)
- R-44 "Nonlinear Excitation in the Ionosphere", A.Y. Wong (March 1969)
- R-45 "Observation of First-Order Ion Energy Distribution in Ion Acoustic Waves, H. Ikezi and R. Taylor (March 1969)
- R-46 "A New Representative for the Conductivity Tensor of a Collisionless Plasma in a Magnetic Field", B.D. Fried and C. Hedrick (March 1969)
- R-47 "Direct Measurements of Linear Growth Rates and Nonlinear Saturation Coefficients", F. Hai (April 1969)
- R-48 "Electron Precipitation Pulsations", F. Coroniti and C.F. Kennel (April 1969)
- R-49 "Auroral Micropulsation Instability", F. Coroniti and C.F. Kennel (May 1969)
- R-50 "Effect of Folker-Plank Collisions on Plasma Wave Echoes", G. Johnston (June 1969)
- R-51 "Linear and Nonlinear Theory of Grid Excitation of Low Frequency Waves in a Plasma", G. Johnston (July 1969)
- R-52 "Theory of Stability of Large Amplitude Periodic (BGK) Waves in Collisionless Plasmas", M.V. Goldman (June 1969)
- R-53 "Observation of Strong Ion Wave-Wave Interaction", R. Taylor and H. Ikezi (August 1969)
- R-55 "Optical Mixing in a Magnetoactive Plasma", G. Weyl (August 1969)
- R-56 "Trapped Particles and Echoes", A.Y. Wong and R. Taylor (Oct. 1969)
- R-57 "Formation and Interaction of Ion-Acoustic Solitons", H. Ikezi, R.J. Taylor and D.R. Baker (July 1970)

- R-58 "Observation of Collisionless Electrostatic Shocks", R. Taylor, D. Baker and H. Ikezi (Dec. 1969)
- R-59 "Turbulent Loss of Ring Current Protons", J.M. Cornwall, F.V. Coroniti and R.M. Thorne (Jan. 1970)
- R-60 "Efficient Modulation Coupling Between Electron and Ion Resonances in Magnetoactive Plasmas", A. Wong, D.R. Baker, N. Booth (Dec. 1969)
- R-61 "Interaction of Quasi-Transverse and Quasi-Longitudinal Waves in an Inhomogeneous Vlasov Plasma", C.L. Hedrick (Jan. 1970)
- R-62 "Observation of Strong Ion-Acoustic Wave-Wave Interaction", R.J. Taylor and H. Ikezi (Jan. 1970)
- R-63 "Perturbed Ion Distributions in Ion-Waves and Echoes", H. Ikezi and R. Taylor (Jan. 1970)
- R-64 "Propagation of Ion Cyclotron Harmonic Wave", E.R. Ault and H. Ikezi (Nov. 1970)
- R-65 "The Analytic and Asymptotic Properties of the Plasma Dispersion Function", A. Banos, Jr. and G. Johnston (Feb. 1970)
- R-66 "Effect of Ion-Ion Collision and Ion Wave Turbulence on the Ion Wave Echo", Dan Baker (June 1970)
- R-67 "Dispersion Discontinuities of Strong Collisionless Shocks", F.V. Coroniti (March 1970)
- R-68 "An Ion Cyclotron Instability", E.S. Weibel (April 1970)
- R-69 "Turbulence Structure of Finite-Beta Perpendicular Fast Shocks", F.V. Coroniti (April 1970)
- R-70 "Steepening of Ion Acoustic Waves and Formation of Collisionless Electrostatic Shocks", R. Taylor (April 1970)
- R-71 "A Method of Studying Trapped Particles Behavior in Magnetic Geometries", C.S. Liu and A.Y. Wong (April 1970)
- R-72 "A Note on the Differential Equation  $g'' + x^2 g = 0$ ", E.S. Weibel (April 1970)
- R-73 "Plasma Response to a Step Electric Field Greater than the Critical Runaway Field, With and Without an Externally Applied Magnetic Field", J.E. Robin (June 1970)
- R-74 "The UC Mathematical On-line Systems as a Tool for Teaching Physics", B.D. Fried and R. White (August 1970)
- R-75 "High Frequency Hall Current Instability", K. Lee, C.F. Kennel, J.M. Kindel (August 1970)
- R-76 "Laminar Wave Train Structure of Collisionless Magnetic Slow Shocks", F.V. Coroniti (Sept. 1970)
- R-77 "Field Aligned Current Instabilities in the Topside Ionosphere", J.M. Kindel and C.F. Kennel (Aug. 1970)
- R-78 "Spatial Cyclotron Damping", Craig Olson (September 1970)
- R-79 "Electromagnetic Plasma Wave Propagation Along a Magnetic Field", C.L. Olson (September 1970)
- R-80 "Electron Plasma Waves and Free-Streaming Electron Bursts", H. Ikezi, P.J. Barrett, R.B. White and A.Y. Wong (Nov. 1970)
- R-81 "Relativistic Electron Precipitation During Magnetic Storm Main Phase", R.M. Thorne and C.F. Kennel (Nov. 1970)
- R-82 "A Unified Theory of SAR Arc Formation at the Plasmopause", J.M. Cornwall, F.V. Coroniti and R.M. Thorne (Nov. 1970)
- R-83 "Nonlinear Collisionless Interaction between Electron and Ion Modes in Inhomogeneous Magnetoactive Plasmas", N. Booth (Dec. 1970)
- R-85 "Remote Double Resonance Coupling of Radar Energy to Ionospheric Irregularities", C.F. Kennel (Jan. 1971)
- R-86 "Ion Acoustic Waves in a Multi-ion Plasma", B.D. Fried, R. White, T. Samec (Jan. 1971)
- R-87 "Current-Driven Electrostatic and Electromagnetic Ion Cyclotron Instabilities", D.W. Forslund, C.F. Kennel, J. Kindel (Feb. 1971)